



## Ecological indicators based on quantitative eDNA metabarcoding: the case of marine reserves

Loïc Sanchez<sup>a,b,\*</sup>, Emilie Boulanger<sup>c</sup>, Véronique Arnal<sup>b</sup>, Pierre Boissery<sup>e</sup>, Alicia Dalongeville<sup>a</sup>, Tony Dejean<sup>d</sup>, Julie Deter<sup>f</sup>, Nacim Guellati<sup>a</sup>, Florian Holon<sup>f</sup>, Jean-Baptiste Juhel<sup>a</sup>, Philippe Lenfant<sup>g</sup>, Fabien Leprieur<sup>a</sup>, Alice Valentini<sup>d</sup>, Stéphanie Manel<sup>a</sup>, David Mouillot<sup>a,h</sup>

<sup>a</sup> MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Montpellier, France

<sup>b</sup> CEFE, Univ Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France

<sup>c</sup> Aix-Marseille Université, Université de Toulon, CNRS, IRD, Mediterranean Institute of Oceanography (MIO), UM 110, 13288, Marseille, France

<sup>d</sup> SPYGEN, 73370 Le Bourget-du-Lac, France

<sup>e</sup> Agence de l'Eau Rhône-Méditerranée-Corse, Délégation de Marseille2, rue Barbusse, CS 90464, 13207 Marseille Cedex 01, France

<sup>f</sup> Andromède Océanologie, 7 place Cassan – Carnon plage, 34130 Mauguio, France

<sup>g</sup> Univ. Perpignan Via Domitia, Centre de Formation et de Recherche sur les Environnements Méditerranéens, UMR 5110, 58 Avenue Paul Alduy, F-66860 Perpignan, France

<sup>h</sup> Institut Universitaire de France, Paris, France

### ARTICLE INFO

#### Keywords:

Community ecology  
Quantitative metabarcoding  
eDNA  
Marine reserves

### ABSTRACT

In a context of marine biodiversity erosion, the need to better understand the effects of overfishing stands out. New genetic techniques such as environmental DNA (eDNA) metabarcoding have emerged and allow the detection of a wider range of species compared to conventional methods, but still fall short of providing reliable abundance estimations and subsequent ecological indicators. In this paper, we propose a combination of metabarcoding and quantitative polymerase chain reaction to obtain the quantity of eDNA molecules per species. This method was used inside and outside six no-take Mediterranean marine reserves to measure the effect of the protection on fish species and build a new indicator. Even if the total quantity of fish eDNA molecules was not different between the inside and outside of the reserves, we detected that cryptobenthic fish eDNA was significantly associated to the outside of reserves. Based on this observation, we propose a novel ecological indicator, the Demerso-pelagic to Benthic fish eDNA Ratio (DeBRA), taking advantage of the eDNA capacity to detect cryptobenthic reef fishes which are often missed by classical surveys. The DeBRA was significantly higher inside reserves, reflecting a higher relative quantity of eDNA molecules belonging to pelagic and demersal fishes under protection against fishing, therefore it appears to be a reliable eDNA-based indicator of human pressure. Furthermore, the DeBRA was not sensitive to habitat or environmental variations and does not require a complete reference database of eDNA sequences since it can rely on sequences assigned at the genus or family scale if possible and necessary.

### 1. Introduction

In marine and freshwater ecosystems, fish support a wide range of functions and contribute to nutrient cycles, carbon sequestration and the provision of vital nutrients to human populations (Holmlund and Hammer, 1999; Golden, 2016; Villéger et al., 2017; Mariani et al., 2020; Robinson et al., 2022). Through anthropogenic activities, fish and their habitats are under an ever-increasing number of threats such as overfishing or pollution at the local scale (Lloret et al., 2020; Cheminée et al.,

2021) and climate warming at the global scale (Smale et al., 2019). These severe impacts jeopardize the nature's contributions to people provided by fish (Hicks et al., 2019; McLean et al., 2019; Robinson et al., 2019; Beger, 2021). To counteract these threats, Marine Protected Areas (MPAs) have become the main management tools on coastal ecosystems to maintain key habitats and viable fish populations (Grorud-Colvert et al., 2021). To reinforce conservation efforts in face of unprecedented biodiversity losses (Dulvy et al., 2021), international governing bodies have recently called for a new global target of 30% coverage by MPAs

\* Corresponding author at: MARBEC, 093 Pl. Eugène Bataillon, 34090 Montpellier, France.


E-mail address: [loic.sanchez@umontpellier.fr](mailto:loic.sanchez@umontpellier.fr) (L. Sanchez).

<https://doi.org/10.1016/j.ecolind.2022.108966>


Received 10 January 2022; Received in revised form 10 May 2022; Accepted 11 May 2022

Available online 26 May 2022

1470-160X/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



**Fig. 1.** Map of the 6 sampled regions in the Northwestern Mediterranean Sea. Inside each region, 4 replicates were sampled in 3 different sites, one inside the marine reserve and two outside.




**Fig. 2.** Histogram showing the distribution of the log-transformed quantity of eDNA molecules per transect for all fish species combined (a) and per fish species per transect (b). Values colored in orange correspond to species absences or false negatives. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

before 2030 (Jones et al., 2020).

Beyond coverage, there is a need to both comprehensively understand the effects of overfishing and to measure the ecological effectiveness of MPAs on fish communities. No-take or fully protected MPAs, also called marine reserves (Costello and Ballantine, 2015), have proven their ability to restore fish stocks and to produce more offspring from large individuals (Marshall et al., 2019; Ban et al., 2019). Yet, the effects of marine reserves on fish communities are not fully understood with some limited benefits (Loiseau et al., 2021) and even counter-intuitive results on biodiversity (Boulanger et al., 2021).

Whether they aim to measure management effectiveness or human impacts, various ecological indicators have been proposed (Meehan et al., 2020). However, most ecological indicators are based on surveys conducted with fishing gears, underwater visual censuses (UVCs), remote underwater videos (RUVs) and baited remote underwater videos (BRUVs), which (1) are selective and lead to biased estimates of biodiversity (Whitmarsh et al., 2017; Costello et al., 2017); (2) require highly advanced taxonomic expertise (Thomsen and Willerslev, 2015); or (3) are destructive or invasive (scientific trawling). The bias can be particularly striking when considering small species such as cryptobenthic fishes, hidden in the substrate, that cannot be correctly monitored with most fishing gears, UVCs or BRUVs (Smith-Vaniz et al., 2006; Alzate et al., 2014) while constituting the building blocks of reef ecosystems (Brandl et al., 2019). Large fish like sharks are also missed by most surveys since they are nocturnal, elusive, or rare (Boussarie et al., 2018). So new census methods with associated indicators are urgently needed to better assess the effects of protection, management and exploitation of coastal ecosystems.

Environmental genetic techniques such as environmental DNA (hereafter eDNA) metabarcoding have shown their ability to accurately detect the presence of a broad range of fishes in the marine environment, (Juhel et al., 2020) and elusive species (Boussarie et al., 2018), outperforming many classical techniques for fish inventories (Thomsen et al., 2012; Fediajevaite et al., 2021; Polanco Fernández et al., 2021). Yet, some studies highlight that eDNA provides poor estimates of benthic communities when considering a large variety of taxa and when sampling the water column (Antich et al., 2021; Laroche et al., 2020) but see Nichols et al. (2022). Moreover, detection ability depends on eDNA shedding and degradation rates that may vary within and among species (Thomsen et al., 2012; Sassoubre et al., 2016) but also with



**Fig. 3.** Boxplot of the log-transformed quantities of total fish eDNA molecules per transect inside and outside marine reserves. No difference was detected with a Wilcoxon test ( $W = 553$ ,  $p$ -value = 0.343).

environmental conditions (Harrison et al., 2019). eDNA detection currently relies on polymerase chain reaction (PCR), a process that may provide results biased in favor of certain species (Bellemain et al., 2010; Clarke et al., 2014; Kelly et al., 2019). Furthermore, species abundances, which inform various key indicators of ecosystem status (Stuart-Smith et al., 2013; Cinner et al., 2020), are still poorly estimated using eDNA in the marine environment; but see Stoeckle et al. (2021).

A recent review, based on 12 studies involving the eDNA metabarcoding of freshwater or marine fish species, reported a coefficient of determination ( $R^2$ ), so a predictive power, ranging from 0.30 to 0.98 for fish counts and a  $R^2$  of 0.75 for fish biomass in controlled environments (Rourke et al., 2021). In natural environments, the  $R^2$  ranged from 0.24 to 0.68 for relative abundance, and from 0.26 to 0.45 for biomass. This difference was also reported in a meta-analysis on species-specific surveys, in which eDNA particle quantity explained 82% and 57% of the observed variation in fish abundance in laboratory experiments and natural environments, respectively (Yates et al., 2019). Indeed, many abiotic and biotic factors such as temperature, depth, activity, and metabolic rates render the estimation of species abundance from eDNA metabarcoding challenging (Rourke et al., 2021). Moreover, depending on possible primer mismatches, there is a chance to under- or over-amplify the barcode sequence of some species or families (Kelly et al., 2019), which in turn would bias the relative abundance of eDNA molecules in a sample, after amplification. More research is needed to understand the extent of this bias and understand whether certain species, e.g. conservation-dependent species, could be underestimated by common markers. In general, this potential bias would be conserved throughout all samples and within families so it would not impact between-site comparisons.

In addition to the difficulty of linking the quantity of eDNA to species abundance, the distribution of eDNA molecules is often heterogeneously distributed in the environment (Turner et al., 2014; Turner et al., 2015). So, a large volume of filtered seawater (100 L) over long transects with many replicates may be required to obtain reliable estimates of species richness, but this sampling effort is often unrealistic in terms of deployment or cost (Stauffer et al., 2021). In light of these potential limitations, we need to test the ability of eDNA-based indicators to differentiate protected vs. unprotected marine areas using archetypal situations where fish communities are strongly affected by human pressure.

The Mediterranean Sea is amongst the most impacted marine areas

(Cavan and Hill, 2021) with a strong decline of most commercial fish stocks (Colloca et al., 2013; Vasilakopoulos et al., 2014; Di Minin et al., 2019). MPAs of varying sizes and levels of protection have been implemented (Gorud-Colvert et al., 2021) and multiple studies have demonstrated the benefits of marine reserves on fish biomass (Giakoumi et al., 2017; Blowes et al., 2020). In this study, we conducted real-time quantitative PCR (qPCR) on eDNA samples filtered inside and outside six Mediterranean marine reserves to provide a quantity of eDNA molecules for each species in each sample. We make the hypothesis that the quantity of eDNA molecules is (i) higher inside protected areas owing to higher fish biomass and (ii) influenced by the position of fish in the water column with crypto-benthic species being less affected by fisheries (Boulanger et al. 2021) and less likely to release large eDNA quantities in the open sea than their demersal or pelagic counterparts. To test these hypotheses, we compared the quantity of eDNA molecules per species, and per category of species (i.e., cryptobenthic, benthic, demersal, and pelagic) inside and outside the reserves through two different approaches. First, we used statistical models to test the effect of protection and position in the water column on the quantity of fish eDNA molecules, while taking into account variability in fish body size. Second, we used a non-statistical approach based on Indicator Values to include both species eDNA quantities and occurrences within sites (Cáceres and Legendre, 2009). We then proposed the Demerso-pelagic to Benthic eDNA Ratio (DeBRa), a novel ecological indicator based on quantitative eDNA to measure the effect of protection or human pressure on fish communities, independently of habitat and environmental conditions.

## 2. Material and methods

### 2.1. Sampling protocol

eDNA samples were collected during the same season, in a two-month period (May-June) with peristaltic pumps (1 L/min) one meter below the surface in six Mediterranean regions, each of them containing a fully protected marine reserve with a strict no-take policy known to provide higher fish biomass than fished areas (Sala and Giakoumi, 2018). In each region, two types of sites were sampled on the same day: inside and outside reserves (Fig. 1). We considered that contamination by dispersal between sampling sites, located 5 to 10 km apart, could be neglected given the strength of the currents in the area (1 km/day), eDNA persistence in the marine environment (24–48 h; Collins et al.,

**Table 1**

Results of GLM that tested the effects of protection, species vertical distribution, length, trophic level, and the environment on the species detection probability and the quantity of DNA molecules. The reference for qualitative variables was the “cryptobenthic” category for vertical distribution, and “outside” for protection. We only show variables selected by the AIC criterion in the final models with statistical significance indicated by \*:  $p < 0.05$ ; \*\*:  $p < 0.01$ ; \*\*\*:  $p < 0.001$ .

Response variable	Variables retained (stepAIC)	Coefficients	Pseudo-R <sup>2</sup>		
Total DNA per transect	PC1	-0.07**	0.14		
	PC3	-0.04			
DNA Detection probability per species per transect	Protection (Inside)	-0.01	0.05		
	Age	-0.99***			
	Trophic levelLog (Common length)	-0.60***			
	Vertical	0.77***			
	Distribution	1.20***			
	Benthic	0.31*			
	Demersal	-0.60***			
	Pelagic	0.11***			
	PC2	0.26			
	PC4	0.36			
	Protection: Benthic	0.58**			
	Protection: Demersal				
	Protection: Pelagic				
DNA per species per transect	Protection (Inside)	-0.44***	0.18		
	Age	0.76***			
	Vertical	-			
	Distribution	0.47***			
	Benthic	0.81***			
	Demersal	0.29***			
	PelagicLog (Common length)	-0.44***			
	Trophic level	-0.20***			
	PC1	-0.08***			
	PC2	0.03**			
	PC3	-0.07***			
	Total cryptobenthic fish DNA per transect	PC1		-0.12	0.09
	Total benthic fish DNA per transect	PC3		-0.13	
Protection (Inside)		-0.18			
Total demersal fish DNA per transect	PC1	-0.05	0.13		
	PC2	0.06			
	PC4	0.06			
Total pelagic fish DNA per transect	Age	0.01	0.23		
	PC1	-0.18**			
	PC3	-0.20**			
Total benthic & cryptobenthic fish DNA per transect	Protection (Inside)	-0.017	0.11		
	PC1	-0.05			
	PC2	0.05*			
Total demersal & pelagic fish DNA per transect	Protection (Inside)	-0.17	0.13		
	PC1	-0.05			
	PC2	0.05			

2018) and recent studies suggesting that eDNA signals display a high spatial fidelity (Murakami et al., 2019; West et al., 2020; Monuki et al., 2022; Jensen et al., 2022). Four replicates of 30L of seawater were sampled at each site along four 2 km-long transects using VigiDNA 0.2 µM filters. After filtration, the capsules were drained with air, filled with a CL1 buffer solution to conserve eDNA molecules, and stored at room temperature before eDNA extraction.

## 2.2. eDNA analyses


eDNA extraction, amplification, and sequencing were described in Boulanger et al. (2021). For the quantification of fish eDNA, the samples were amplified using the teleost-specific 12S mitochondrial rRNA primer (teleo) which has been extensively used for fish eDNA metabarcoding (Cantera et al., 2019; Cilleros et al., 2018; Coutant et al., 2021) and performs better (*i.e.* detects more species, with less bias and more specific amplification) than primers based on alternative loci (Collins et al., 2018; Weigand et al., 2019; Polanco Fernández et al., 2021). Then, the qPCR was performed in a final volume of 25 µL, which included 3 µL of DNA, 12.5 µL of SYBR® Green Master Mix (BioRad®), 7.3 µL of ddH<sub>2</sub>O, 0.5 µL of each “teleo” primer (10 mM), 1 µL of human blocking primer (100 µM; Valentini et al., 2016) and 0.2 g/µL of bovine serum albumin (BSA, Roche Diagnostic). Each sample was analyzed in 3 replicates. A dilution series of a synthetic gene of 10<sup>-1</sup> ng/µL, 10<sup>-3</sup> ng/µL, 10<sup>-5</sup> ng/µL, 10<sup>-7</sup> ng/µL (that correspond to 1,13E + 08, 1,13E + 06, 1,13E + 04 and 1,13E + 02 eDNA molecules respectively) was used as a standard for the qPCR analysis. The tubes containing the eDNA samples were sealed and the qPCR standards were added to the qPCR plate in a room separate from the eDNA extraction room. The qPCR cycle was as follows: 95 °C for 10 min, followed by 55 cycles of 95 °C for 30 s and 55 °C for 30 s. Melting curves were produced by plotting fluorescence intensity against temperature as the temperature was increased from 65 to 95 °C at a rate of 0.5 °C every 5 s. Samples were analyzed on a BIO-RAD® CFX96 Touch real-time PCR detection.

## 2.3. Incidence data

The OBITools package (Boyer et al., 2016) was used to analyze sequence reads following the protocol described in Pont et al. (2018). The taxonomic assignment of reads was performed using the ecotag program with both an in-house local reference database and the sequences retrieved from the European Nucleotide Archive (ENA) database (Boulanger et al., 2021). The in-house database contains 320 fish species, corresponding to 41% of all teleost species in the Mediterranean Sea and to 75% of the regional species pool (Boulanger et al., 2021). Reads showing 98% or more similarity with the reference database were kept. Taxa were preferentially assigned based on the local database, except if the similarity was higher for the ENA database. Considering the incorrect assignment of a few sequences to the sample due to tag-jumps (Schnell et al., 2015), we discarded all sequences with a frequency of occurrence < 0.001 per sequence and per library. The resulting dataset was manually checked to correct erroneous identifications and to remove foreign species and assignments that could not be assigned at the species level (Boulanger et al., 2021). The metabarcoding method provided a quantity of reads per species, *i.e.*, the number of sequences that were detected for each species. Yet, this standardized quantity cannot be compared among samples because the initial eDNA quantity in each sample was yet unknown, but it allows to estimate species proportions in each sample.

## 2.4. eDNA quantity per species

We used a qPCR to obtain the total quantity of fish eDNA molecules in each sample, all species combined. This quantity was corrected in proportion to the quantity of reads that were excluded from the corresponding samples (foreign species, human contamination, unassigned taxa). For each sample, the corrected quantity of total molecules was then multiplied by species proportions in the corresponding sample obtained from the read metabarcoding outputs to calculate the quantity of eDNA molecules per species per sample. In more details, species proportions were obtained by dividing the number of metabarcoding reads of each species by the total number of reads before the bioinformatic analysis in each sample. Samples for which the qPCR did not provide a satisfying total quantity of eDNA (<10,000 molecules) were



**Fig. 4.** Partial regression plots from the binomial GLM model using the *visreg* function (package “visreg”) showing the species detection probability in function of the protection level while controlling for all other fixed effects, for each position in the water column. Error bars represent 95% confidence intervals.

removed from the dataset.

### 2.5. Environmental variables

To describe the habitat and environmental variation among sites, we used the first four axes derived from a Principal Component analysis (PCA) (Boulanger et al., 2021). The PCA was computed using a set of habitat and environmental variables extracted for each site: habitat coverage, distance to land, sea surface temperatures (SST), mean bottom depths, and mean benthic and surface chlorophyll *a*. The PCA was applied to avoid multicollinearity between these variables and model overfitting (Fig. S1). The first four principal components explained 74% of the total variance in environmental conditions.

### 2.6. Modelling the quantity of eDNA

The quantity of DNA molecules per species and per transect was log-transformed ( $\log_{10}(x + 1)$ ) before analyses. Our dataset at the species level contained a large proportion of zeros corresponding to either species absence or false negatives. We thus used a hurdle model, combining two subsequent models whose outputs were analyzed separately: a binomial generalized linear model (GLM) to predict species presence-absence and a gaussian GLM to predict the quantity of DNA molecules per species after excluding all zeros (Liu et al., 2019; Brown et al., 2021). We built these models to test the influence of protection and species categories on total quantity of eDNA molecules per transect

(all species combined) and the total quantity of DNA molecules per transect per category of species: pelagic, demersal, benthic, and cryptobenthic species, as well as demerso-pelagic species (grouping demersal and pelagic species) and benthic species as a combination of cryptobenthic and benthic species. These categories were defined according to the vertical distribution of species in the water column, with pelagic species swimming in the water column, demersal species swimming near the bottom, benthic species living on the bottom and cryptobenthic species being smaller benthic species that mostly hide in the substrate (Brandl et al., 2018). Indeed, species with varying traits according to their morphological profiles (Friedman et al., 2020), and thus varying metabolic activities (Killen et al., 2010) may shed varying quantities of eDNA. Therefore, our models at the species level included the level of protection and fish vertical distribution but also species traits extracted from FishBase such as common length and trophic level to control for other sources of variability among species.

Because the age and size of the reserves ranged from 6 to 44 years of implementation, and from 65 to 1074 ha, we also included the age and size of the reserves. We normalized the age and size and the reserves such that the oldest or biggest value was 1, while fished areas were 0, following the path of Cinner et al. (2018). All models included the four first principal components of the previously mentioned PCA to control for habitat and environmental variability among sites and regions. To avoid collinearity between variables in the models, we used the variance inflation factor (VIF) to confirm that these variables could all be included in the model. We then performed a descending and ascending




Fig. 5. Partial regression plots from the gaussian GLM model using the *visreg* function (package “visreg”), showing the species quantity of DNA molecules in function of the protection level while controlling for all other fixed effects, for each position in the water column. Error bars represent 95% confidence intervals.

*stepAIC* (package MASS) to select the most parsimonious model to avoid overfitting and coefficient biases. Model fit and residuals were visually checked.

### 2.7. Indicator values and effect sizes

As a complementary approach to statistical models and to avoid the issue of zero-inflated distributions, we computed the Indicator Value index *IndVal* (Cáceres and Legendre, 2009) using the function *multipatt* (package *indicpecies*). *IndVal* combines two factors *A* and *B*, respectively corresponding to proxies of species specificity and fidelity for a given group of sites:

$$IndVal_{ij} = \sqrt{A_{ij} \times B_{ij}} \times 100$$

$A_{ij}$  is defined as the mean abundance of the *i*th species within its *j*th protection level divided by the sum of mean abundances of this species over all protection levels, and  $B_{ij}$  is defined as the number of sites where the species occurs inside its target protection level (Cáceres and Legendre, 2009). In other words, a species with 1 in specificity for protected areas will only be found in marine reserves, and a species with 1 in fidelity will be found in each reserve.

*IndVal* is, by construction, ranged between 0 and 100. The maximum value means that the species is only found in its target protection level and is always found in sites pertaining to this protection level. The minimum value means that the species is not found in sites of this protection level. Indicator values were computed for each species in


protected ( $IndVal_i^P$ ) and unprotected ( $IndVal_i^U$ ) areas based on the quantity of eDNA. The following formula was then used to calculate the Effect Size of the reserves for the *i*th species:

$$Effect\ Size_i = (IndVal_i^P - IndVal_i^U) / Max(IndVal_i^P - IndVal_i^U)$$

Because this method cannot account for environmental variability, we selected sites with similar habitat and environmental conditions by plotting the first three axes of the PCA and visually selecting a cluster containing both protected and unprotected sites with similar environments. The cluster represented sites displaying a value lower than  $-0.5$  on PC1 and comprised 8 protected sites and 28 unprotected sites. The effect sizes were computed on both the complete dataset and the cluster of environmentally similar sites. To test whether the Effect Size was significantly influenced by the species category while considering phylogenetic distances between species, we computed a Phylogenetic Generalized Least Squares model (pGLS) with the *ppls* function (package “caper”). The phylogeny used for this model was obtained via the Fish Tree of Life API (Rabosky et al., 2018).

### 2.8. The Demerso-pelagic to Benthic eDNA Ratio (DeBRa)

Considering that benthic species were recently found to be more present outside than inside Mediterranean reserves (Boulangier et al., 2021) and that the most threatened species, which are mainly demersal and pelagic, were found more present inside reserves (Loiseau et al., 2021), we computed the ratio between the log-transformed quantity of



**Fig. 6.** Effect sizes of protection on the detected species computed from the Indicator Values on environmentally similar sites. Only the highest and lowest 25 species were plotted to facilitate reading. Colors indicate the vertical distribution. All species are represented in Fig. S2.

benthic fish eDNA molecules and the log-transformed quantity of demerso-pelagic fish DNA molecules to build a new indicator:

$$DeBRa = \ln(TotaleDNA_{Dem,Pel}) / \ln(TotaleDNA_{Benth})$$

The DeBRa was computed for each site, and the influence of protection was investigated using both a non-parametric rank test, and a gaussian GLM taking into account habitat and environmental variability.

### 3. Results

#### 3.1. Total quantity of eDNA per transect

The total quantity of fish eDNA molecules detected per transect ranged from 6,384 to 1,978,396 and followed a log-normal distribution (Fig. 2a). The Wilcoxon test did not detect a significant difference between inside and outside marine reserves ( $W = 553, p\text{-value} = 0.343$ ) (Fig. 3). The gaussian GLM revealed no effect of protection on the total quantity of eDNA molecules per transect but a strong and significant effect of environmental PC1 (Table 1).

#### 3.2. Quantity of eDNA per species

The quantity of eDNA molecules per species per transect ranged from 0 to 1,965,527 with 94 different species detected throughout all samples

(Fig. S2, Table S1). Zero values represented 48% of the data, while positive values followed a log-normal distribution (Fig. 2b).

The binomial and gaussian GLM models revealed quite similar results, with significant effects of position in the water column and environmental PCA variables (Table 1). The probability to detect a species was similar inside than outside reserves (Fig. 4) while the quantity of eDNA molecules per detected species was markedly higher outside than inside reserves whatever the species category (Fig. 5). Regarding position in the water column, pelagic and cryptobenthic species had a lower detection probability (Fig. 4) and a lower quantity of eDNA molecules (Fig. 5) than benthic and demersal species. However, a significant interaction was found between protection and position in the water column in the binomial model. Pelagic species were more frequently detected inside marine reserves. The chance to be detected inside or outside reserve was equal for cryptobenthic species, and slightly higher inside reserves for benthic and demersal species, albeit non-significant. Furthermore, the gaussian model revealed that lower quantities of eDNA were detected for species with higher trophic level and larger body size. Reserve age had a positive effect on the quantity of eDNA but a negative effect on the detection probability.

#### 3.3. Total quantity of eDNA per species category

The quantity of eDNA molecules per fish species category ranged from 0 to 1,967,876 and followed a log-normal distribution. Among all




Fig. 7. Partial regression plot from the GLM model using the *visreg* function (package “visreg”), showing the Effect Size of the reserve on the quantity of eDNA molecules for different species categories while controlling for other traits (length and trophic levels) and phylogenetic distances.




Fig. 8. Partial regression plot from GLM models using the *visreg* function (package “visreg”), showing the values of the new DeBRa indicator (Demersal to Benthic Ratio) as a function of the protection level while controlling for environmental variability. The model explains 15.9% of the variance.

the models computed for each species category, none revealed a significant effect of protection on the quantity of eDNA (Table 1). After combining benthic with cryptobenthic species and demersal with

pelagic species, no significant effect of protection was detected either. We only detected a strong effect of principal components corresponding to habitat and environmental variables in each of these models

(Table 1).

### 3.4. Indicator values and effect size

Indicator values ranged from 0 to 73.41 in reserves, and from 0 to 71.75 in fished areas, with uniform distributions for both. The Effect Size computed with these values ranged from  $-0.50$  to  $0.64$  and followed a gaussian distribution. The top 5% species with positive Effect Sizes are demersal or pelagic, while the top 5% species with negative values are cryptobenthic and benthic (Fig. S3). In the analysis restricted to a cluster of environmentally similar sites, the same pattern emerged: most positive values (higher inside reserves) correspond to demersal or pelagic species while most negative values (higher outside reserves) correspond to cryptobenthic or benthic species (Fig. 6). The pGLS successfully detected this pattern and indicated increasing Effect Sizes along the water column with significant or nearly significant effects of species category (Fig. 7).

### 3.5. Demerso-pelagic to Benthic eDNA Ratio (DeBRa)

Values for the DeBRa ranged from 1.0197 to 2.9410 and followed a normal distribution. The gaussian GLM revealed a significantly higher ratio of eDNA shed by demerso-pelagic fish than benthic fish in marine reserves compared to fished areas ( $p$ -value = 0.001). No significant effect PC2 ( $p$ -value = 0.1119) was detected, while PC1, PC3 and PC4 were not retained in the model by the stepAIC (Fig. 8). Our model displayed a pseudo- $R^2$  of 0.16. This significant difference between both groups was also detected with a non-parametric rank test of Wilcoxon ( $p$ -value = 0.003), indicating a  $p$ -value more than one order of magnitude smaller than when taking only species richness into account ( $p$ -value = 0.05).

## 4. Discussion

Surprisingly, the GLM predicting the total quantity of eDNA molecules per transect did not reveal the expected effect of protection, but only a significant effect of environmental variables, while fully protected areas or marine reserves are notoriously known to promote higher fish abundance and biomass (Lester et al., 2009; Soykan and Lewison, 2015; Sala and Giakoumi, 2018). This result indicates that the total quantity of fish eDNA molecules cannot reliably be used as a proxy of fish biomass or as an indicator of management or fishing intensity.

At the species level, statistical models revealed a protection effect on the quantity of eDNA molecules. Even when accounting for environmental variability, which had a significant effect, there were more fish eDNA molecules outside than inside MPAs. Until recently, studies had only reported an increase in the abundance of pelagic over demersal species inside reserves (Libralato et al., 2010), but since classical surveys cannot offer such a species coverage for cryptobenthic species there are no previous records of increase in cryptobenthic abundance outside reserves. This finding could be explained by an hypoallometric relationship: a higher number of young and juvenile individuals may shed a higher eDNA quantity per gram of fish biomass than their adult counterparts (Maruyama et al., 2014; Takeuchi et al., 2019), and therefore provide a false signal of higher biomass, while decreasing the chances to amplify species with very low quantities of eDNA, which would explain higher quantities but lower detection rates outside marine reserves for some categories. Yet, our deep sequencing effort (1 million sequences) may partly mitigate this effect. Marine protected areas cover a relatively small area in the Mediterranean (6.81%) and considerable efforts have been made to create artificial nurseries far from predators (e.g., ports), which means that many coastal fish nurseries are outside MPAs (MedPAN et al., 2016; Claudet et al., 2021). The large number of small individuals in these nurseries could explain higher quantities of eDNA molecules outside reserves. This result underlines the current complementarity of eDNA metabarcoding with conventional methods such as visual, acoustic or camera surveys that remain necessary to estimate fish

body size.

The low quantities of eDNA molecules for pelagic species could be explained by their behavior since they often live in schools, are highly mobile and may have shed few and aggregated eDNA molecules in an area where they just passed. The differences in detection and in eDNA quantity between fishes with different positions in the water column could then partly be due to the variability in the shedding capacity between these categories, or simply to metabolic differences. This result is challenging to interpret and suggests that many hypotheses remain to be tested, notably in terms of biological trait influence on eDNA shedding. Finally, since (i) the substantial quantity of zeros demands to decompose our analysis in two subsequent models, each of which is missing a part of reality, (ii) our statistical models display a very low pseudo- $R^2$  and (iii) that no interaction could be detected in the gaussian model between protection level and position in the water column we suggest that working at the species level may not be an appropriate choice if the goal is to develop an ecological indicator based on the quantity of eDNA molecules. Yet, the protection effect was not significant either when computing models on each species category while the effect of environment was always significant as shown by Loiseau et al. (2021).

As an alternative, the Indicator Value approach, which is not based on statistical distributions and assumptions, has the advantage of using all the zeros in the dataset since it relies on both specificity and fidelity measures. In our case, after computing the Effect Sizes for each species, we observed a clear underlying pattern in the data: species with the highest effect size of protection were demersal or pelagic while those with the lowest effect size of protection were benthic or cryptobenthic. The pGLS further confirmed that the differences in effect size between species categories were significant while guaranteeing they were not due to phylogenetic similarities between species or other traits. Using Indicator Values we thus detected a marked pattern that could not be detected with species-based models.

Fishing seems to decrease the ratio between demerso-pelagic and benthic species eDNA quantity, meaning that fishing may favor benthic species to the detriment of demersal and pelagic species (Boulanger et al., 2021). In our study, the increase of the DeBRa indicator inside reserves indicates an increase in the number of demerso-pelagic fish eDNA molecules compared to benthic fish eDNA molecules, independently of habitat and environmental variables making this indicator transferable and comparable between ecosystems and seasons. Yet, this indicator does not provide information at the population level: we cannot determine if this increase is due to a simple increase in the density of adults inside reserves, or if it was caused by a massive increase in benthic juveniles outside during the spring season. Indeed, this hypothesis should be considered as many benthic species have pelagic early life stages, that could be less predated by demersal species, themselves under fishing pressure. Current research is exploring ways to use eDNA and environmental RNA (eRNA) to determine life-history stages from genetic material (Yates et al., 2021), so these methods could contribute to better understand whether unprotected areas may or not play an important and complementary role of nursery for some species.

## 5. Conclusion

Our findings suggest that eDNA metabarcoding coupled to quantitative real-time PCR can be used in field surveys to quantify eDNA molecules pertaining to each detected species. We show that Indicator Values could be used to detect some patterns that would not be detected by classical statistical models based on either species occurrences or abundances given the high proportion of zeros inherent to eDNA surveys. We also show that we cannot base our indicator on the total quantity of eDNA in a transect, and that the differences in quantities lie at the species level, with effects of many variables affecting the detected quantity of eDNA, such as body size, trophic level, environmental variables but also age of the reserve. More generally, our results emphasize

the difficulty to evaluate benefits of MPAs on fish communities. We reveal antagonistic responses to protection between species categories, highlighting the need to base management strategies on multiple indicators and not only on the biomass of some targeted fishes. In the Mediterranean Sea, and more precisely in the Northwest regional pool, more than 75% of species are referenced in our genetic database, but in tropical regions which are species richness hotspots (Tittensor et al., 2010), a large proportion of species are missing in genetic reference databases (Marques et al., 2021). Since the DeBRA indicator only requires to determine if the species is cryptobenthic, benthic, demersal, or pelagic, it could be possible to assign unknown sequences at the genus or family level (for genera and families that have 95% of their taxa pertaining to the same category, for example), and thus confidently build this indicator without assignation at the species level. We expect that this new indicator can provide consistent results in other temperate and tropical regions to reveal human impacts or management benefits regardless of habitat, seasonal and environmental conditions.

### CRedit authorship contribution statement

**Loïc Sanchez:** Conceptualization, Investigation, Formal analysis, Visualization, Writing – original draft, Writing – review & editing. **Emilie Boulanger:** Investigation, Writing – review & editing. **Véronique Arnal:** Writing – review & editing. **Pierre Boissery:** Writing – review & editing. **Alicia Dalongeville:** Conceptualization, Formal analysis, Visualization, Writing – review & editing. **Tony Dejean:** Methodology, Writing – review & editing. **Julie Deter:** Investigation, Writing – review & editing. **Nacim Guellati:** Investigation, Writing – review & editing. **Florian Holon:** Investigation, Writing – review & editing. **Jean-Baptiste Juhel:** Investigation, Writing – review & editing. **Philippe Lenfant:** Investigation, Writing – review & editing. **Fabien Leprieur:** Formal analysis, Writing – review & editing. **Alice Valentini:** Methodology, Software, Writing – review & editing. **Stéphanie Manel:** Conceptualization, Methodology, Funding acquisition, Writing – review & editing. **David Mouillot:** Conceptualization, Formal analysis, Writing – original draft, Funding acquisition, Writing – review & editing

### Funding

Funding was provided by the Agence de l'Eau RMC, the Region Occitanie and the RESERVEBENEFIT project through the 2015–2016 BiodivERSA COFUND call with the ANR (France).

### Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Stephanie Manel reports financial support was provided by Rhone Mediterranee Corsica Region Water Agency. Stephanie Manel reports financial support was provided by French National Research Agency. Stephanie Manel reports financial support was provided by Occitanie Region.

### Acknowledgements

The authors thank the funders of the study, the managers of the studied MPAs, and we are grateful to Seguin R. and Loiseau N. for their technical support and comments on an earlier version of this article.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ecolind.2022.108966>.

### References

- Alzate, A., Zapata, F.A., Giraldo, A., 2014. A comparison of visual and collection-based methods for assessing community structure of coral reef fishes in the Tropical Eastern Pacific. *RBT* 62, 359. <https://doi.org/10.15517/rbt.v62i0.16361>.
- Antich, A., Palacín, C., Cebrian, E., Golo, R., Wangensteen, O.S., Turon, X., 2021. Marine biomonitoring with eDNA: Can metabarcoding of water samples cut it as a tool for surveying benthic communities? *Mol. Ecol.* 30, 3175–3188. <https://doi.org/10.1111/mec.15641>.
- Ban, N.C., Gurney, G.G., Marshall, N.A., Whitney, C.K., Mills, M., Gelcich, S., Bennett, N. J., Meehan, M.C., Butler, C., Ban, S., Tran, T.C., Cox, M.E., Breslow, S.J., 2019. Well-being outcomes of marine protected areas. *Nat. Sustain.* 2, 524–532. <https://doi.org/10.1038/s41893-019-0306-2>.
- Bellemain, E., Carlsen, T., Brochmann, C., Coissac, E., Taberlet, P., Kauserud, H., 2010. ITS as an environmental DNA barcode for fungi: an *in silico* approach reveals potential PCR biases 9. <https://doi.org/10.1186/1471-2180-10-189>.
- Beger, M., 2021. Accepting the loss of habitat specialists in a changing world. *Nat. Ecol. Evol.* 5, 556–557. <https://doi.org/10.1038/s41559-021-01402-6>.
- Blowes, S.A., Chase, J.M., Di Franco, A., Frid, O., Gotelli, N.J., Guidetti, P., Knight, T.M., May, F., McGlenn, D.J., Micheli, F., Sala, E., Belmaker, J., 2020. Mediterranean marine protected areas have higher biodiversity via increased evenness, not abundance. *J. Appl. Ecol.* 57, 578–589. <https://doi.org/10.1111/1365-2664.13549>.
- Boulanger, E., Loiseau, N., Valentini, A., Arnal, V., Boissery, P., Dejean, T., Deter, J., Guellati, N., Holon, F., Juhel, J.-B., Lenfant, P., Manel, S., Mouillot, D., 2021. Environmental DNA metabarcoding reveals and unpacks a biodiversity conservation paradox in Mediterranean marine reserves, Dryad, Dataset. *Proc. Royal Soc. B* 288 (20210112), 1–10. <https://doi.org/10.1098/rspb.2021.0112>.
- Boussarie, G., Bakker, J., Wangensteen, O.S., Mariani, S., Bonnin, L., Juhel, J.-B., Kiszka, J.J., Kulbicki, M., Manel, S., Robbins, W.D., Vigliola, L., Mouillot, D., 2018. Environmental DNA illuminates the dark diversity of sharks. *Sci. Adv.* 4, eaap9661. <https://doi.org/10.1126/sciadv.aap9661>.
- Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., Coissac, E., 2016. *obtools*: a UNIX-inspired software package for DNA metabarcoding. *Mol. Ecol. Resour.* 16, 176–182. <https://doi.org/10.1111/1755-0998.12428>.
- Brandl, S.J., Goatley, C.H.R., Bellwood, D.R., Tornabene, L., 2018. The hidden half: ecology and evolution of cryptobenthic fishes on coral reefs. *Biol. Rev.* 93, 1846–1873. <https://doi.org/10.1111/brv.12423>.
- Brandl, S.J., Tornabene, L., Goatley, C.H.R., Casey, J.M., Morais, R.A., Côté, I.M., Baldwin, C.C., Parravicini, V., Schiethkatte, N.M.D., Bellwood, D.R., 2019. Demographic dynamics of the smallest marine vertebrates fuel coral reef ecosystem functioning. *Science* 364, 1189–1192. <https://doi.org/10.1126/science.aav3384>.
- Brown, C.J., Mellin, C., Edgar, G.J., Campbell, M.D., Stuart-Smith, R.D., 2021. Direct and indirect effects of heatwaves on a coral reef fishery. *Glob. Change Biol.* 27, 1214–1225. <https://doi.org/10.1111/gcb.15472>.
- Cáceres, M.D., Legendre, P., 2009. Associations between species and groups of sites: indices and statistical inference. *Ecology* 90, 3566–3574. <https://doi.org/10.1890/08-1823.1>.
- Cantera, I., Cilleros, K., Valentini, A., Cerdan, A., Dejean, T., Iribar, A., Taberlet, P., Vigouroux, R., Brosse, S., 2019. Optimizing environmental DNA sampling effort for fish inventories in tropical streams and rivers. *Sci. Rep.* 9, 3085. <https://doi.org/10.1038/s41598-019-39399-5>.
- Cavan, E.L., Hill, S.L., 2021. Commercial fishery disturbance of the global ocean biological carbon sink. *Glob. Change Biol.* 00, 1–10. <https://doi.org/10.1111/gcb.16019>.
- Cheminée, A., Le Direach, L., Rouanet, E., Astruch, P., Goujard, A., Blanfuné, A., Bonhomme, D., Chassaing, L., Jouvenel, J.Y., Ruitton, S., Thibaut, T., Harmelin-Vivien, M., 2021. All shallow coastal habitats matter as nurseries for Mediterranean juvenile fish. *Sci. Rep.* 11, 14631. <https://doi.org/10.1038/s41598-021-93557-2>.
- Cilleros, K., Valentini, A., Allard, L., Dejean, T., Etienne, R., Grenouillet, G., Iribar, A., Taberlet, P., Vigouroux, R., Brosse, S., 2018. Unlocking biodiversity and conservation studies in high diversity environments using environmental DNA (eDNA): A test with Guianese freshwater fishes. *Mol. Ecol. Resour.* 19, 27–46. <https://doi.org/10.1111/1755-0998.12900>.
- Cinner, J., Maire, E., Huchery, C., Mouillot, D., 2018. Gravity of human impacts mediates coral reef conservation gains. *PNAS* 115, 27. <https://doi.org/10.1073/pnas.1708001115>.
- Cinner, J.E., Zamborain-Mason, J., Gurney, G.G., Graham, N.A.J., MacNeil, M.A., Hoey, A.S., Mora, C., Villéger, S., Maire, E., McClanahan, T.R., Maina, J.M., Kittinger, J.N., Hicks, C.C., D'agata, S., Huchery, C., Barnes, M.L., Feary, D.A., Williams, I.D., Kulbicki, M., Vigliola, L., Wantiez, L., Edgar, G.J., Stuart-Smith, R.D., Sandin, S.A., Green, A.L., Beger, M., Friedlander, A.M., Wilson, S.K., Brokovich, E., Brooks, A.J., Cruz-Motta, J.J., Booth, D.J., Chabanet, P., Tupper, M., Ferse, S.C.A., Sumaila, U.R., Hardt, M.J., Mouillot, D., 2020. Meeting fisheries, ecosystem function, and biodiversity goals in a human-dominated world. *Science* 368, 307–311. <https://doi.org/10.1126/science.aax9412>.
- Coutant, O., Cantera, I., Cilleros, K., Dejean, T., Valentini, A., Murielle, J., Brosse, S., 2021. Detecting fish assemblages with environmental DNA: Does protocol matter? Testing eDNA metabarcoding method robustness. *Environ. DNA* 3, 619–630.
- Clarke, L.J., Soubrier, J., Weyrich, L.S., Cooper, A., 2014. Environmental metabarcodes for insects: *in silico* PCR reveals potential for taxonomic bias. *Mol. Ecol. Resour.* 14, 1160–1170. <https://doi.org/10.1111/1755-0998.12265>.
- Claudet, J., Loiseau, C., Pebayle, A., 2021. Critical gaps in the protection of the second largest exclusive economic zone in the world. *Marine Policy* 124, 104379. <https://doi.org/10.1016/j.marpol.2020.104379>.

- Collins, R.A., Wangenstein, O.S., O'Gorman, E.J., Mariana, S., Sims, D.W., Genner, M.J., 2018. Persistence of environmental DNA in marine systems. *Commun. Biol.* 1, 185. <https://doi.org/10.1038/s42003-018-0192-6>.
- Colloca, F., Cardinale, M., Maynou, F., Giannoulaki, M., Scarella, G., Jenko, K., Bellido, J.M., Fiorentino, F., 2013. Rebuilding Mediterranean fisheries: a new paradigm for ecological sustainability: Sustainability of Mediterranean fisheries. *Fish. Fish.* 14, 89–109. <https://doi.org/10.1111/j.1467-2979.2011.00453.x>.
- Costello, M.J., Ballantine, B., 2015. Biodiversity conservation should focus on no-take Marine Reserves: 94% of Marine Protected Areas allow fishing. *Trends Ecol. Evol.* 30 (9), 507–509. <https://doi.org/10.1016/j.tree.2015.06.011>.
- Costello, M. J., Basher, Z., McLeod, L., Asaad, I., Claus, S., Vandepitte, L., Yasuhara, M., Gislason, H., Edwards, M., Appeltans, Enevoldsen, H., Edgar, G. J., Miloslavich, P., De Monte, S., Sousa Pinto, I., Obura, D., Bates, A. E., 2017. Methods for the study of Marine Biodiversity. In: *The GEO Handbook on Biodiversity Observation Networks*. [https://doi.org/10.1007/978-3-319-27288-7\\_6](https://doi.org/10.1007/978-3-319-27288-7_6).
- Di Minin, E., Brooks, T.M., Toivonen, T., Butchard, S.H.M., Heikinheimo, V., Watson, J.E. M., Burgess, N.D., Challender, D.W.S., Goettsch, B., Jenkins, R., Moilanen, A., 2019. Identifying global centers of unsustainable commercial harvesting of species. *Sci. Adv.* 5 (4) <https://doi.org/10.1126/sciadv.aau2879>.
- Dulvy, N.K., Pacoureau, N., Rigby, C.L., Pollom, R.A., Jabado, R.W., Ebert, D.A., Finucci, B., Pollock, C.M., Cheok, J., Derrick, D.H., Herman, K.B., Sherman, C.S., VanderWright, W.J., Lawson, J.M., Walls, R.H.L., Carlson, J.K., Charvet, P., Bineesh, K.K., Fernando, D., Ralph, G.M., Matsushiba, J.H., Hilton-Taylor, C., Fordham, S.V., Simpfendorfer, C.A., 2021. Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Curr. Biol.* 31 (21) <https://doi.org/10.1016/j.cub.2021.08.062>.
- Fediajevaite, J., Priestley, V., Arnold, R., Savolainen, V., 2021. Meta-analysis shows that environmental DNA outperforms traditional surveys, but warrants better reporting standards. *Ecol. Evol.* 11, 4803–4815. <https://doi.org/10.1002/ece3.7382>.
- Friedman, S.T., Price, S.A., Corn, K.A., Larouche, O., Martinez, C.M., Wainwright, P.C., 2020. Body shape diversification along the benthic–pelagic axis in marine fishes. *Proc. R. Soc. B* 287, 20201053. <https://doi.org/10.1098/rspb.2020.1053>.
- Giakoumi, S., Scianna, C., Plass-Johnson, J., Micheli, F., Grorud-Colvert, K., Thiriet, P., Claudet, J., Di Carlo, G., Di Franco, A., Gaines, S.D., Garcia-Charton, J.A., Lubchenco, J., Reimer, J., Sala, E., Guidetti, P., 2017. Ecological effects of full and partial protection in the crowded Mediterranean Sea: a regional meta-analysis. *Sci. Rep.* 7, 8940. <https://doi.org/10.1038/s41598-017-08850-w>.
- Golden, C., 2016. Fall in fish catch threatens human health. *Nature* 534, 317–320.
- Grorud-Colvert, K., et al., 2021. The MPA guide: a framework to achieve global goals for the ocean. *Science* 373 (6560). <https://doi.org/10.1126/science.abc0861>.
- Harrison, J.B., Sunday, J.M., Rogers, S.M., 2019. Predicting the fate of eDNA in the environment and implications for studying biodiversity. *Proc. R. Soc. B* 286, 20191409. <https://doi.org/10.1098/rspb.2019.1409>.
- Hicks, C.C., Cohen, P.J., Graham, N.A.J., Nash, K.L., Allison, E.H., D'Lima, C., Mills, D.J., Roscher, M., Thilsted, S.H., Thorne-Lyman, A.L., MacNeil, M.A., 2019. Harnessing global fisheries to tackle micronutrient deficiencies. *Nature* 574, 95–98. <https://doi.org/10.1038/s41586-019-1592-6>.
- Holmlund, C.M., Hammer, M., 1999. Ecosystem services generated by fish populations. *Ecol. Econ.* 29, 253–268. [https://doi.org/10.1016/S0921-8009\(99\)00015-4](https://doi.org/10.1016/S0921-8009(99)00015-4).
- Jensen, M.R., Sigsgaard, E.E., Avila, M.D.P., Agersnap, S., Brenner-Larsen, P., Sengupta, M.E., Xing, Y., Krag, M.A., Knudsen, S.W., Carl, H., Møller, P.R., Thomsen, P.F., 2022. Short-term temporal variation of coastal marine eDNA. *Environ. DNA* 00, 1–16. <https://doi.org/10.1002/edn3.285>.
- Jones, K.R., Klein, C.J., Grantham, H.S., Possingham, H.P., Halpern, B.S., Burgess, N.D., Butchart, S.H.M., Robinson, J.G., Kingston, N., Bhola, N., Watson, J.E.M., 2020. Area requirements to safeguard Earth's marine species. *One Earth* 2 (2), 188–196. <https://doi.org/10.1016/j.oneear.2020.01.010>.
- Juhel, J.-B., Utama, R.S., Marques, V., Vimono, I.B., Sugeha, H.Y., Pouyaud, L., Dejean, T., Mouillot, D., Hocdé, R., 2020. Accumulation curves of environmental DNA sequences predict coastal fish diversity in the coral triangle. *Proc. R. Soc. B* 287, 20200248. <https://doi.org/10.1098/rspb.2020.0248>.
- Kelly, R.P., Shelton, A.O., Gallego, R., 2019. Understanding PCR Processes to Draw Meaningful Conclusions from Environmental DNA Studies. *Sci. Rep.* 9, 12133. <https://doi.org/10.1038/s41598-019-48546-x>.
- Killen, S.S., Atkinson, D., Glazier, D.S., 2010. The intraspecific scaling of metabolic rate with body mass in fishes depends on lifestyle and temperature. *Ecol. Lett.* 13, 184–193. <https://doi.org/10.1111/j.1461-0248.2009.01415.x>.
- Laroche, O., Kersten, O., Smith, C.R., Goetze, E., 2020. From sea surface to seafloor: a benthic allochthonous eDNA survey for the abyssal ocean. *Front. Mar. Sci.* 7 <https://doi.org/10.3389/fmars.2020.00682>.
- Lester, S., Halpern, B., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B., Gaines, S., Airamé, S., Warner, R., 2009. Biological effects within no-take marine reserves: a global synthesis. *Mar. Ecol. Prog. Ser.* 384, 33–46. <https://doi.org/10.3354/meps08029>.
- Libralato, S., Coll, M., Tempesta, M., Santojanni, A., Spoto, M., Palomera, I., Arneri, E., Solidoro, C., 2010. Food-web traits of protected and exploited areas of the Adriatic Sea. *Biol. Conserv.* 143, 2182–2194. <https://doi.org/10.1016/j.biocon.2010.06.002>.
- Liu, L., Shih, Y.-C.-T., Strawderman, R.L., Zhang, D., Johnson, B.A., Chai, H., 2019. Statistical Analysis of Zero-Inflated Nonnegative Continuous Data: A Review. *Statist. Sci.* 34 <https://doi.org/10.1214/18-ST681>.
- Lloret, J., Biton-Porsmoguer, S., Carreño, A., Di Franco, A., Sahyoun, R., Melià, P., Claudet, J., Sève, C., Ligas, A., Belharet, M., Calò, A., Carbonara, P., Coll, M., Corrales, X., Lembo, G., Sartor, P., Bitetto, I., Vilas, D., Piroddi, C., Prato, G., Charbonnel, E., Bretton, O., Hartmann, V., Prats, L., Font, T., 2020. Recreational and small-scale fisheries may pose a threat to vulnerable species in coastal and offshore waters of the western Mediterranean. *ICES J. Mar. Sci.* 77, 2255–2264. <https://doi.org/10.1093/icesjms/fsz071>.
- Loiseau, N., Thuiller, W., Stuart-Smith, R.D., Devicor, V., Edgar, G.J., Velez, L., Cinner, J.E., Graham, N.A.J., Renaud, J., Hoey, A.S., Manel, S., Mouillot, D., 2021. Maximizing regional biodiversity requires a mosaic of protection levels. *PLoS Biol.* 19, e3001195. <https://doi.org/10.1371/journal.pbio.3001195>.
- Mariani, G., Cheung, W.W.L., Lyet, A., Sala, E., Mayorga, J., Velez, L., Gaines, S.D., Dejean, T., Troussellier, M., Mouillot, D., 2020. Let more big fish sink: Fisheries prevent blue carbon sequestration—half in unprofitable areas. *Sci. Adv.* 6, eabb4848. <https://doi.org/10.1126/sciadv.abb4848>.
- Marques, V., Castagné, P., Fernández, A.P., Borrero-Pérez, G.H., Hocdé, R., Guérin, P.-É., Juhel, J.-B., Velez, L., Loiseau, N., Letessier, T.B., Bessudo, S., Valentini, A., Dejean, T., Mouillot, D., Pellissier, L., Villéger, S., 2021. Use of environmental DNA in assessment of fish functional and phylogenetic diversity. *Conserv. Biol.* 35, 1944–1956. <https://doi.org/10.1111/cobi.13802>.
- Marshall, D.J., Gaines, S., Warner, R., Barneche, D.R., Bode, M., 2019. Underestimating the benefits of marine protected areas for the replenishment of fished populations. *Front. Ecol. Environ.* 17, 407–413. <https://doi.org/10.1002/fee.2075>.
- Maruyama, A., Nakamura, K., Yamanaka, H., Kondoh, M., Minamoto, T., 2014. The Release Rate of Environmental DNA from Juvenile and Adult Fish. *PLoS ONE* 9, e114639. <https://doi.org/10.1371/journal.pone.0114639>.
- McLean, M.J., Mouillot, D., Goascoz, N., Schlaich, I., Auber, A., 2019. Functional reorganization of marine fish nurseries under climate warming. *Glob. Change Biol.* 25, 660–674. <https://doi.org/10.1111/gcb.14501>.
- MedPAN et al., 2016. Le Statut des Aires Marines Protégées de Méditerranée en 2016 : Résultats principaux. Brochure MedPAN & ONU Environnement - PAM - CAR/ASP.
- Meehan, M.C., Ban, N.C., Devillers, R., Singh, G.G., Claudet, J., 2020. How far have we come? A review of MPA network performance indicators in reaching qualitative elements of Aichi Target 11. *Conserv. Lett.* 13 <https://doi.org/10.1111/conl.12746>.
- Monuki, K., Barber, P.H., Gold, Z., 2022. eDNA captures depth partitioning in a kelp forest ecosystem. *PLoS ONE* 16 (11), e0253104. <https://doi.org/10.1371/journal.pone.0253104>.
- Murakami, H., Yoon, S., Kasai, A., Minamoto, T., Yamamoto, S., Sakata, M.K., Horiuchi, T., Sawada, H., Kondoh, M., Yamashita, Y., Masuda, R., 2019. Dispersion and degradation of environmental DNA from caged fish in a marine environment. *Fish. Sci.* 85, 327–337. <https://doi.org/10.1007/s12562-018-1282-6>.
- Nichols, P.K., Timmers, M., Marko, P.B., 2022. Hide 'n' seek: Direct versus indirect metabarcoding of coral reef cryptic communities. *Environ. DNA* 4, 93–107. <https://doi.org/10.1002/edn3.203>.
- Polanco Fernández, A., Marques, V., Fopp, F., Juhel, J., Borrero-Pérez, G.H., Cheutin, M., Dejean, T., González Corredor, J.D., Acosta-Chaparro, A., Hocdé, R., Eme, D., Maire, E., Spescha, M., Valentini, A., Manel, S., Mouillot, D., Albouy, C., Pellissier, L., 2021. Comparing environmental DNA metabarcoding and underwater visual census to monitor tropical reef fishes. *Environ. DNA* 3, 142–156. <https://doi.org/10.1002/edn3.140>.
- Pont, D., Rocle, M., Valentini, A., Civade, R., Jean, P., Maire, A., Roset, N., Schabuss, M., Zornig, H., Dejean, T., 2018. Environmental DNA reveals quantitative patterns of fish biodiversity in large rivers despite its downstream transportation. *Sci. Rep.* 8, 10361. <https://doi.org/10.1038/s41598-018-28424-8>.
- Rabosky, D.L., Chang, J., Title, P.O., Cowman, P.F., Sallan, L., Friedman, M., Kaschner, K., Garilao, C., Near, T.J., Coll, M., Alfaro, M.E., 2018. An inverse latitudinal gradient in speciation rate for marine fishes. *Nature* 559, 392–395. <https://doi.org/10.1038/s41586-018-0273-1>.
- Robinson, J.P.W., Wilson, S.K., Jennings, S., Graham, N.A.J., 2019. Thermal stress induces persistently altered coral reef fish assemblages. *Glob. Change Biol.* 25, 2739–2750. <https://doi.org/10.1111/gcb.14704>.
- Robinson, J.P.W., Maire, E., Bodin, N., Hempson, T.N., Graham, N.A.J., Wilson, S.K., MacNeil, M.A., Hicks, C., 2022. Climate-induced increases in micronutrient availability for coral reef fisheries. *One Earth.* <https://doi.org/10.1016/j.oneear.2021.12.005>.
- Rourke, M.L., Fowler, A.M., Hughes, J.M., Broadhurst, M.K., DiBattista, J.D., Fielder, S., Wilkes Walburn, J., Furlan, E.M., 2021. Environmental DNA (eDNA) as a tool for assessing fish biomass: A review of approaches and future considerations for resource surveys. *Environmental DNA* edn3.185. <https://doi.org/10.1002/edn3.185>.
- Sala, E., Giakoumi, S., 2018. No-take marine reserves are the most effective protected areas in the ocean. *ICES J. Mar. Sci.* 75, 1166–1168. <https://doi.org/10.1093/icesjms/fox059>.
- Sassoubre, L.M., Yamahara, K.M., Gardner, L.D., Block, B.A., Boehm, A.B., 2016. Quantification of Environmental DNA (eDNA) Shedding and Decay Rates for Three Marine Fish. *Environ. Sci. Technol.* 50, 10456–10464. <https://doi.org/10.1021/acs.est.6b03114>.
- Schnell, I.B., Bohmann, K., Gilbert, M.T.P., 2015. Tag jumps illuminated – reducing sequence-to-sample misidentifications in metabarcoding studies. *Mol. Ecol. Resour.* 15, 1289–1303. <https://doi.org/10.1111/1755-0998.12402>.
- Smale, D.A., Wernberg, T., Oliver, E.C.J., Thomsen, M., Harvey, B.P., Straub, S.C., Burrows, M.T., Alexander, L.V., Benthushy, J.A., Donat, M.G., Feng, M., Hobday, A. J., Holbrook, N.J., Perkins-Kirkpatrick, S.E., Scannell, H.A., Sen Gupta, A., Payne, B. L., Moore, P.J., 2019. Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nat. Clim. Chang.* 9, 306–312. <https://doi.org/10.1038/s41558-019-0412-1>.
- Smith-Vaniz, W.F., Jelks, H.L., Rocha, L.A., 2006. Relevance of cryptic fishes in biodiversity assessments: a case study at Buck Island Reef National Monument St. Croix. *Bull. Marine Sci.* 79, 32.

- Soykan, C.U., Lewison, R.L., 2015. Using community-level metrics to monitor the effects of marine protected areas on biodiversity: Biodiversity Metrics for MPAs. *Conserv. Biol.* 29, 775–783. <https://doi.org/10.1111/cobi.12445>.
- Stauffer, S., Jucker, M., Keggins, T., Marques, V., Andrello, M., Bessudo, S., Cheutin, M.-C., Borrero-Pérez, G.H., Richards, E., Dejean, T., Hocdé, R., Juhel, J.-B., Ladino, F., Letessier, T.B., Loiseau, N., Maire, E., Mouillot, D., Martinezguerra, M.M., Manel, S., Fernández, A.P., Valentini, A., Velez, L., Albouy, C., Pellissier, L., Waldoock, C., 2021. How many replicates to accurately estimate fish biodiversity using environmental DNA on coral reefs? (preprint). *Ecology*. <https://doi.org/10.1101/2021.05.26.445742>.
- Stoeckle, M.Y., Adolf, J., Charlop-Powers, Z., Dunton, K.J., Hinks, G., VanMorter, S.M., 2021. Trawl and eDNA assessment of marine fish diversity, seasonality, and relative abundance in coastal New Jersey, USA. *ICES J. Mar. Sci.* 78 (1), 293–304. <https://doi.org/10.1093/icesjms/fsaa225>.
- Stuart-Smith, R.D., Bates, A.E., Lefcheck, J.S., Duffy, J.E., Baker, S.C., Thomson, R.J., Stuart-Smith, J.F., Hill, N.A., Kininmonth, S.J., Airoidi, L., Becerro, M.A., Campbell, S.J., Dawson, T.P., Navarrete, S.A., Soler, G.A., Strain, E.M.A., Willis, T.J., Edgar, G.J., 2013. Integrating abundance and functional traits reveals new global hotspots of fish diversity. *Nature* 501, 539–542. <https://doi.org/10.1038/nature12529>.
- Takeuchi, A., Iijima, T., Kakuzen, W., Watanabe, S., Yamada, Y., Okamura, A., Horie, N., Mikawa, N., Miller, M.J., Kojima, T., Tsukamoto, K., 2019. Release of eDNA by different life history stages and during spawning activities of laboratory-reared Japanese eels for interpretation of oceanic survey data. *Sci. Rep.* 9, 6074. <https://doi.org/10.1038/s41598-019-42641-9>.
- Thomsen, P.F., Kielgast, J., Iversen, L.L., Møller, P.R., Rasmussen, M., Willerslev, E., 2012. Detection of a Diverse Marine Fish Fauna Using Environmental DNA from Seawater Samples. *PLoS ONE* 7, e41732. <https://doi.org/10.1371/journal.pone.0041732>.
- Thomsen, P.F., Willerslev, E., 2015. Environmental DNA – An emerging tool in conservation for monitoring past and present biodiversity. *Biol. Conserv.* 183, 4–18. <https://doi.org/10.1016/j.biocon.2014.11.019>.
- Tittensor, D.P., Mora, C., Jetz, W., Lotze, H.K., Ricard, D., Berghe, E.V., Worm, B., 2010. Global patterns and predictors of marine biodiversity across taxa. *Nature* 466, 1098–1101. <https://doi.org/10.1038/nature09329>.
- Turner, C.R., Barnes, M.A., Xu, C.C.Y., Jones, S.E., Jerde, C.L., Lodge, D.M., 2014. Particle size distribution and optimal capture of aqueous microbial eDNA. *Methods Ecol. Evol.* 5, 676–684. <https://doi.org/10.1111/2041-210X.12206>.
- Turner, C.R., Uy, K.L., Everhart, R.C., 2015. Fish environmental DNA is more concentrated in aquatic sediments than surface water. *Biol. Conserv.* 183, 93–102. <https://doi.org/10.1016/j.biocon.2014.11.017>.
- Valentini, A., Taberlet, P., Miaud, C., Cívade, R., Herder, J., Thomsen, P.F., Bellemain, E., Besnard, A., Coissac, E., Boyer, F., Gaboriaud, C., Jean, P., Poulet, N., Roset, N., Copp, G.H., Geniez, P., Pont, D., Argillier, C., Baudoin, J.-M., Peroux, T., Crivelli, A. J., Olivier, A., Acqueberge, M., Le Brun, M., Møller, P.R., Willerslev, E., Dejean, T., 2016. Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Mol. Ecol.* 25, 929–942. <https://doi.org/10.1111/mec.13428>.
- Vasilakopoulos, P., Maravelias, C.D., Tserpes, G., 2014. The Alarming Decline of Mediterranean Fish Stocks. *Curr. Biol.* 24, 1643–1648. <https://doi.org/10.1016/j.cub.2014.05.070>.
- Villéger, S., Brosse, S., Mouchet, M., Mouillot, D., Vanni, M.J., 2017. Functional ecology of fish: current approaches and future challenges. *Aquat. Sci.* 79, 783–801. <https://doi.org/10.1007/s00027-017-0546-z>.
- Weigand, H., Beermann, A.J., Ciampor, F., Costa, F.O., Csabai, Z., Duarte, S., Geiger, M. F., Grabowski, M., Rimet, F., Rulik, B., Strand, M., Szucsich, N., Weigand, A.M., Willassen, E., Wyler, S.A., Bouchez, A., Borja, A., Ciamporová-Zatovičová, Z., Ferreira, S., Dijkstra Klaas-Douwé, K.D.B., Eisendle, U., Freyhof, J., Gadawski, P., Graf, W., Haegerbaeumer, A., van der Hoorn, B.B., Japoshvili, B., Keresztes, L., Keskin, E., Leese, F., Macher, J.N., Mamos, T., Paz, G., Pešić, V., Pfannkuchen, D.M., Ferreira, S., Dijkstra Klaas-Douwé, K.D.B., Eisendle, U., Freyhof, J., Gadawski, P., Ekrm, T., 2019. DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-analysis and recommendations for future work. *Sci. Total Environ.* 678, 499–524. <https://doi.org/10.1016/j.scitotenv.2019.04.247>.
- West, K.M., Stat, M., Harvey, E.S., Skepper, C.L., DiBattista, J.D., Richards, Z.T., Travers, M.J., Newman, S.J., Bunce, M., 2020. eDNA metabarcoding survey reveals fine-scale coral reef community variation across a remote, tropical island ecosystem. *Mol. Ecol.* 29, 1069–1086. <https://doi.org/10.1111/mec.15382>.
- Whitmarsh, S.K., Fairweather, P.G., Huvneers, C., 2017. What is Big BRUVver up to? Methods and uses of baited underwater video. *Rev. Fish Biol. Fish.* 27, 53–73. <https://doi.org/10.1007/s11160-016-9450-1>.
- Yates, M.C., Derry, A.M., Cristescu, M.E., 2021. Environmental RNA: A Revolution in Ecological Resolution? *Trends Ecol. Evol.* 36, 601–609. <https://doi.org/10.1016/j.tree.2021.03.001>.
- Yates, M.C., Fraser, D.J., Derry, A.M., 2019. Meta-analysis supports further refinement of eDNA for monitoring aquatic species-specific abundance in nature. *Environ. DNA* 1, 5–13. <https://doi.org/10.1002/edn3.7>.